

Omex Folicao

FUNCTION

Omex Folicao is a high concentration liquid foliar fertiliser, designed to be applied to the foliar parts of the cocoa tree and many other crops.

Regular applications of Omex Folicao will improve the quality and yield of cocoa and other crops.

Omex Folicao has a very positive impact on the performance of the cocoa, with more abundant and larger pods and improved quality cocoa beans.

DESCRIPTION

A fully water soluble suspension fertiliser containing NPK, magnesium and chelated trace elements.



Analysis of Omex Folicao

			Wt/Wt*	Wt/Vol
Nitrogen		N	8.70%	13.50%
Nitrate			6.60%	10.30%
Ammonium			2.10%	3.20%
Phosphate		P ₂ O ₅	17.20%	27.00%
Potassium		K ₂ O	17.20%	27.00%
Magnesium		MgO	1.00%	1.60%
Iron	(EDTA)	Fe	0.106%	0.166%
Zinc	(EDTA)	Zn	0.053%	0.083%
Copper	(EDTA)	Cu	0.053%	0.083%
Manganese	(EDTA)	Mn	0.053%	0.083%
Boron		B	0.022%	0.034%
Cobalt	(EDTA)	Co	0.0008%	0.0012%
Molybdenum		Mo	0.0008%	0.0012%
pH (10% solution)			4.0-5.0	
Specific Gravity			1.54-1.58 @ 18°C	

*For registration purposes Wt/Wt concentrations are recommended.

DIRECTIONS FOR USE

The spray tank should be filled with half of the required amount of water. Measure the required amount of Omex Folicao and add to the tank whilst maintaining constant agitation.

Add remaining water to correct dilution. Spray.



OMEX
AGRIFLUIDS

www.omex.com

DILUTION: Recommended water rate is 500-1500 Litres per hectare

Always shake container before opening



Recommendations for use:

CROP	RATE l/ha	RATE ml/100l	APPLICATION DETAILS
Cocoa	2.0-4.0	200-400 (30-60ml/15l)	Apply during periods of active growth, if deficiency severe repeat applications at 10-14 days
Beans	2.0-4.0	200-400	Apply during periods of active growth, if deficiency severe repeat applications at 10-14 days
Brassicas	2.0-4.0	200-400	Apply during periods of active growth, if deficiency severe repeat applications at 10-14 days
Maize, Rice, Sorghum	2.0-4.0	200-400	Apply during periods of active growth, if deficiency severe repeat applications at 10-14 days
Potatoes	2.0-4.0	200-400	Apply during periods of active growth, if deficiency severe repeat applications at 10-14 days
Peas	2.0-4.0	200-400	Apply during periods of active growth, if deficiency severe repeat applications at 10-14 days
Tomatoes	2.0-4.0	200-400	Apply during periods of active growth, if deficiency severe repeat applications at 10-14 days
Cassava	2.0-4.0	200-400	Apply during periods of active growth, if deficiency severe repeat applications at 10-14 days

TANK MIXING COMPATIBILITY

Omex Folicao is compatible with most, but not all, pesticides, growth regulators and micro-nutrients with regard to physical tank mixing and biological effects on the crop. However, Omex cannot accept any liability for any loss or damage as not all pesticides have been tested and because the efficacy of any mix will depend on, among other factors, the pesticide concerned, crop conditions, growth stage, weather and volumes of water used.

PRECAUTIONS

Omex Folicao should be stored in frost free conditions with optimum storage range between 5-40°C. Omex Folicao is a non-hazardous and non-flammable foliar fertiliser. Gloves and face shield should be worn when handling the concentrate. In situations of prolonged storage there may be slight settling of the nutrient particles. This is reversible on shaking.

PACKING: Omex Folicao is available in HDPE drums of 0.25L, 0.5L, 1L, 3L, 5L, 10L capacity. Drums are fitted with tamper evident closures. 200 and 1000 Litre mini bulk available.

Omex Agrifluids Limited

Saddlebow Road • King's Lynn • Norfolk PE34 3JA • UK

t: +44 (0)1553 817 500 • f: +44 (0)1553 817 501

e: agrifluids@omex.com • www.omex.com